

# Lateral™

## Soil Wetting Agent

**CONTAINS NON-PLANT FOOD INGREDIENTS**

**ACTIVE INGREDIENTS:**

90% nonionic polyoxyethylene-polyoxypropylene block copolymer

10% alpha-(para-Nonylphenyl)-omega-hydroxypoly(oxyethylene)

**NOT RECOMMENDED FOR USE AS A FERTILIZER SUBSTITUTE**

## KEEP OUT OF REACH OF CHILDREN

Read label before use. Keep container tightly closed. Keep only in original container. Avoid breathing spray. Wash hands thoroughly after handling. Do not eat, drink or smoke when using this product. Wear protective gloves, protective clothing, and eye protection.

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfco.org/metals.htm>

**NET CONTENTS: 2.5 U.S. Gallons (9.46 L)**

**DENSITY: 8.70 lbs/gal at 68 deg. F**

**NET WEIGHT: 21.75 lbs at 68 deg. F**

**EXACTO**   
Better products, real solutions  
200 Old Factory Road, Sharon, WI 53585  
262-854-9040

## FIRST AID

*Have the product container with you when calling a poison control center or doctor, or going for treatment.*

**IF IN EYES:** Rinse cautiously with water for several minutes. Remove contact lenses if present and easy to do then continue rinsing.

**IF ON SKIN OR CLOTHING:** Take off contaminated clothing and wash it before reuse.

**IF SWALLOWED:** Unless advised otherwise by a poison control center or doctor, have person rinse mouth with water.

**IF INHALED:** Remove victim to fresh air and keep at rest in a position comfortable for breathing.

## GENERAL INFORMATION

**Lateral™** is a soil wetting agent concentrate that improves moisture uniformity and wetting in hydrophobic soils that are susceptible to Localized Dry Spot (LDS).

## DIRECTIONS FOR USE

**NOTE: Lateral is a soil water management system and is NOT effective on foliage.**

When using low volume spray equipment and an over-the-top application, treatment must be immediately followed with a minimum ¼ inch of water to ensure soil contact.

Recommended for all turf grass areas including golf course tees, fairways, athletic fields, lawns, sod farms, and any other general use turf grass areas.

**Lateral** treatments may be made approximately once a month.

The first treatment is made at a high use rate in order to properly surround the water repellent soil or thatch layers with surfactant.

Treatments that follow replace the surfactant lost over time due to normal degradation and biological activity under average conditions.

Treatments should start prior to or when signs of visible water stress begin to show, or if water is observed pooling on the surface.

WARRANTY STATEMENT: EXACTO®, INC. (Exacto) represents and warrants that its products are manufactured in accordance with the product's specifications and that its products, at the time of delivery to the purchaser, shall be free from all material defects. EXACTO MAKES NO WARRANTIES OF ANY KIND, EXPRESS OR IMPLIED, CONCERNING THE PRODUCT, INCLUDING BUT NOT LIMITED TO THE IMPLIED WARRANTIES OF MERCHANTABILITY AND FITNESS FOR A PARTICULAR PURPOSE, ARE DISCLAIMED. FURTHER, IT IS UNDERSTOOD AND AGREED THAT EXACTO'S LIABILITY SHALL NOT EXCEED THE RETURN OF THE AMOUNT OF THE PURCHASE PRICE PAID BY PURCHASER AND UNDER NO CIRCUMSTANCES SHALL EXACTO BE LIABLE FOR SPECIAL, INDIRECT, INCIDENTAL, CONSEQUENTIAL OR PUNITIVE DAMAGES. Products for which a warranty claim is made are to be returned transportation prepaid to Exacto's production facility. Any use not in accordance with product directions and precautionary statements, and any modifications or additives to the products, or any adulteration, dilution or alteration by the purchaser shall void this warranty. No employee or representative of Exacto is authorized to change this warranty in any way or to grant any other warranty.

**Lateral** may be applied via an in-ground irrigation system, through a spray bar on a tractor or via hand-held sprayers.

**Lateral** may be applied to bare ground, stressed or healthy plants if adequate water is used in the application.

Do not pre-mix with pesticides, nutrients or other adjuvants before mixing with water.

If the user does not have experience with the mixture being applied, perform a jar test using the order and amounts in the mixture and perform a phytotoxicity test to ensure crop safety.

## USE RATES

**First application:** 4 to 6 ounces of **Lateral** mixed in a minimum of 2 gallons of water for every 1000 square feet of turf or soil to be treated.

**Second, third, fourth, etc. application:** 2 to 4 ounces of **Lateral** in a minimum of 2 gallons of water for every 1000 square feet of turf or soil that will be treated.

Applications should be made every 30 days.

## STORAGE AND DISPOSAL

**Storage:** Store in a cool well-ventilated place. Keep in original container tightly closed. Do not reuse empty container. Do not store with food, feed, or other material to be used or consumed by humans or animals. Do not contaminate water supplies. For optimal storage, store between 40° and 90° F.

**Disposal:** Do not contaminate water, food or feed by storage or disposal. Dispose of contents in container to an approved waste disposal facility in accordance with all local, regional, national and international regulations. Triple rinse (or equivalent) adding rinse water to application tank. Offer container for recycling or dispose of in a sanitary landfill or by other procedures approved by local regulations.